

**SOLUTIONS & ANSWERS FOR KERALA ENGINEERING  
ENTRANCE EXAMINATION-2011  
VERSION – A1**

**[PHYSICS & CHEMISTRY]**

1. Ans: 6%

$$\begin{aligned} \text{Sol: } \frac{\Delta p}{p} &= \frac{\Delta m}{m} + \frac{\Delta V}{V} \\ &= \frac{0.05}{5} \times 100 + \frac{0.05}{1} \times 100 \\ &= 6\% \end{aligned}$$

2. Ans:  $\text{MLT}^{-1}$  and  $\text{MLT}^{-4}$

$$\begin{aligned} \text{Sol: According to principle of homogeneity} \\ [\text{MLT}^{-2}] &= a\text{T}^{-1} \\ \therefore a &= [\text{MLT}^{-1}] \\ [\text{MLT}^{-2}] &= b\text{T}^2 \\ \therefore b &= [\text{MLT}^{-4}] \end{aligned}$$

3. Ans:  $30 \text{ km h}^{-1}$

$$\begin{aligned} \text{Sol: Average velocity} &= 40 \text{ km h}^{-1} \\ \frac{2v_1 v_2}{v_1 + v_2} &= 40 \\ \frac{2 \times 60 \times v_2}{60 + v_2} &= 40 \\ \text{Solving, } v_2 &= 30 \text{ km h}^{-1} \end{aligned}$$

4. Ans: 8 s


$$\begin{aligned} \text{Sol: } 48 + \frac{1}{2}at^2 &= 10t \\ 48 + \frac{1}{2}t^2 &= 10 \\ \text{Solving, } t &= 8 \text{ s} \end{aligned}$$

5. Ans:  $\sqrt{\frac{u^2 + v^2}{2}}$

Sol: Standard results.

6. Ans:  $196.3 \text{ m s}^{-1}$

Sol:



$$\begin{aligned} \tan 30 &= \frac{10v}{3400} \\ v &= \frac{340}{\sqrt{3}} = 196.3 \text{ m s}^{-1} \end{aligned}$$

7. Ans:  $30^\circ$

$$\begin{aligned} \text{Sol: } H_1 &= H_2 \\ u_1^2 \sin^2 45 &= u_2^2 \sin^2 \theta \\ \sin^2 \theta_2 &= \frac{u_1^2}{u_2^2} \cdot \sin^2 45 \\ &= \frac{1}{2} \cdot \frac{1}{2} \\ \sin \theta &= \frac{1}{2} \Rightarrow \theta = 30^\circ \end{aligned}$$

8. Ans:  $\sqrt{\frac{a}{2b}}$

$$\begin{aligned} \text{Sol: } y &= bx^2 \\ \Rightarrow \frac{dy}{dt} &= 2bx \frac{dx}{dt} \quad \text{---(i)} \\ \frac{dy}{dt} &= at \quad (\Theta v_y = u_y + a_y t) \\ \Rightarrow at &= 2bx \frac{dx}{dt} \\ \Rightarrow at dt &= 2bx dx \\ \Rightarrow \int at dt &= \int 2bx dx \\ \frac{at^2}{2} &= bx^2 + c \quad \text{---(ii)} \\ \text{At, } t &= 0, x = 0 \Rightarrow c = 0 \\ \text{(ii)} \Rightarrow \frac{at^2}{2} &= bx^2 \\ \Rightarrow x &= \sqrt{\frac{at^2}{2b}} = \sqrt{\frac{a}{2b}} t \\ \therefore v_x &= \frac{dx}{dt} = \sqrt{\frac{a}{2b}} \end{aligned}$$

9. Ans:  $40 \text{ m s}^{-1}$


$$\begin{aligned} \text{Sol: } P_x &= 2 \times 8 = 16 \\ P_y &= 1 \times 12 = 12 \end{aligned}$$

Momentum of third piece =  $p = \sqrt{16^2 + 12^2} = 20$

Velocity =  $\frac{p}{m} = \frac{20}{0.5} = 40 \text{ m s}^{-1}$

10. Ans: 2 : 1

Sol:



$$L = \frac{1}{2} g \cos 60 t_1^2$$

$$L \cos \theta = \frac{1}{2} g t_2^2$$

$$\frac{t_1^2}{t_2^2} = \frac{1}{\cos^2 60} = 4$$

$$t_1 : t_2 = 2 : 1$$

11. Ans:  $20 \text{ m s}^{-1}$

$$\text{Sol: } v = \sqrt{gr} = \sqrt{10 \times 40} = 20 \text{ m s}^{-1}$$

12. Ans:  $\frac{2mg(h+x)}{x^2}$

$$\text{Sol: } mg(h+x) = \frac{1}{2} kx^2$$

Solving

$$k = \frac{2mg(h+x)}{x^2}$$

13. Ans: 25

$$\text{Sol: } \frac{mg(2-1.5)}{mg \cdot 2} \times 100 = 25\%$$

14. Ans: The potential energy of the particle is zero.

Sol: In horizontal plane PE remains constant equal to zero, assuming surface to be the zero level.

15. Ans:  $\sqrt{gh}$

$$\text{Sol: } v^2 = \frac{2gh}{1 + \frac{k^2}{r^2}} \text{ for ring } k^2 = r^2$$

$$= \frac{2gh}{2} = gh$$

$$v = \sqrt{gh}$$

16. Ans:  $\frac{L}{4}$

$$\text{Sol: } L^2 = 2KI = 2K \frac{L}{\omega}$$

$$L = \frac{2K}{\omega}$$

$$L' = \frac{2 \left( \frac{K}{2} \right)}{2\omega} = \frac{L}{4}$$

17. Ans:  $\frac{20}{9} \text{ m}$

$$\text{Sol: } a = \left( \frac{m_2 - m_1}{m_1 + m_2} \right) g = \frac{10}{3}$$

$$S = \frac{1}{2} at^2 = \frac{20}{3}$$

$$M_X = \frac{2 \times \frac{20}{3} - 1 \times \frac{20}{3}}{3} = \frac{20}{9}$$

18. Ans: 1.36%

$$\text{Sol: } B = \frac{P}{\frac{\Delta V}{V}}$$

$$\frac{\Delta V}{V} = \frac{P}{B} = \frac{\rho gh}{B}$$

Substituting = 1.36%

19. Ans: 1 : 8

$$\text{Sol: } T^2 \propto R^3$$

$$\frac{T_1^2}{T_2^2} = \left( \frac{R}{4R} \right)^3$$

$$\frac{T_1}{T_2} = \frac{1}{8}$$

20. Ans:  $\sqrt{3} \times 11.2 \text{ km s}^{-1}$

$$\text{Sol: } KE = \frac{1}{2} mv^2 - \frac{1}{2} m \times (11.2)^2$$

$$= \frac{1}{2} m (2 \times 11.2)^2 - \frac{1}{2} m \times (11.2)^2$$

$$\frac{1}{2} m v^2 = 3 \times \frac{1}{2} m \times 11.2^2$$

$$V = \sqrt{3} \times 11.2 \text{ km s}^{-1}$$

21. Ans:  $0.1 \text{ m s}^{-1}$

$$\text{Sol: } v = \frac{2}{9} \frac{(\rho - \sigma) r^2 g}{\eta} \propto (\rho - \sigma)$$

$$\frac{v_1}{v_2} = \frac{\rho_1 - \sigma}{\rho_2 - \sigma} = 0.1$$

22. Ans:  $\frac{a}{\sqrt{2\pi}}$

Sol:  $v^2 = \rho g x$   
 $a^2 \sqrt{\rho g x} = \pi r^2 \sqrt{\rho g x} \times 2$   
 $r = \frac{a}{\sqrt{2\pi}}$

23. Ans: Liquid in B increases.

Sol: Let M gram of ice is floating in liquid of density 1.2. Its displaced volume is  
 $\frac{m \text{ c.c.}}{1.2} < m \text{ c.c.}$

When it melts it occupies m c.c

24. Ans: Alloys have larger values of Young's modulus than metals.

Sol: Knowledge based.

25. Ans: 600 K

Sol:  $\frac{T_1 - T_2}{T_1} = 0.4$   
 $T_1 - T_2 = 0.4 T_1$   
 $T_2 = 0.6 T_1$   
 $\frac{T_1' - T_2}{T_1'} = 0.5$   
 $T_1' = \frac{0.6}{0.5} T_1 = 600 \text{ K}$

26. Ans:  $\frac{7}{5}$

Sol:  $C_P = \frac{7}{2} R$   
 $C_V = \frac{5}{2} R$   
 $\therefore r = \frac{C_P}{C_V} = \frac{7}{5}$

27. Ans: Isochoric process

Sol:  $W = 0$   
 $\therefore dV = 0$   
 $\therefore V = \text{constant}$

28. Ans: 80 °C

Sol:  $\frac{1}{2} \times \frac{1}{2} m v^2 = \frac{1}{4} m \times 4 \times 10^4$   
 $= 125 \times \Delta T \times m$   
 $\Delta T = \frac{4 \times 10^4}{500} = 80 \text{ }^\circ\text{C}$

29. Ans: 6.25 cm

Sol:  $T = 2\pi \sqrt{\frac{m}{K}}$   
 $mg = Kx$   
 $\text{Solving } x = 6.25 \text{ cm}$

30. Ans:  $3^3$

Sol:  $A = A_0 e^{\frac{-bt}{2m}}$   
 $\therefore \text{Amplitude becomes } \frac{1}{27} \text{ times after 6 seconds}$

31. Ans: 1 : 2

Sol:  $mg = 2 K x_A = K x_B$   
 $\frac{x_A}{x_B} = \frac{1}{2}$   
 $\frac{W_A}{W_B} = \frac{F x_A}{F x_B} = \frac{1}{2}$

32. Ans: 2 : 1

Sol:  $\frac{v_0}{v_c} = \frac{\frac{v}{2\lambda}}{\frac{v}{4\lambda}} = 2$

33. Ans:  $\sqrt{\frac{P}{\rho}}$

Sol:  $c = \sqrt{\frac{\gamma P}{\rho}}$

34. Ans: No correct answer. Data is inconsistent.

35. Ans:  $\frac{100Q}{\epsilon_0}$

Sol: Charge per metre = 100Q  
 $\phi = \frac{1}{\epsilon_0} q = \frac{100Q}{\epsilon_0}$

36. Ans:  $\frac{r}{4}$



$$\frac{KQq}{r} = \frac{1}{2} mv^2$$

$$\frac{1}{2} m \cdot 4v^2 = \frac{KQq}{r}$$

$$4 = \frac{r}{r'} \Rightarrow r' = \frac{r}{4}$$

**37.** Ans:  $\pi$

Sol:  $U = -pE \cos\theta$   
For  $U$  to maximum  
 $\cos\theta = -1 \Rightarrow \theta = \pi$

**38.** Ans:  $\frac{1}{3} \times 10^{-9}$  N

Sol:  $\frac{KQ_1}{3} = 10, \frac{KQ_2}{1} = 10$   
 $KQ_1 = 30 \times 10^{-2}, KQ_2 = 10 \times 10^{-2}$   
$$\frac{KQ_1 Q_2}{10^{-2}} = \frac{(30 \times 10^{-2})(10 \times 10^{-2})}{K \times 10^{-2}}$$
  
 $F = \frac{1}{3} \times 10^{-9}$  N

**39.** Ans: 0.5

Sol: Standard result.

**40.** Ans: 15.6  $\Omega$

Sol:  $R = \frac{\rho\lambda}{A}$   
 $R \propto \lambda^2$  for given volume  
$$\frac{R_1}{R_2} = \frac{\lambda_1^2}{\lambda_2^2}$$
  
 $R_2 = 15.6 \Omega$

**41.** Ans:  $2 \times 10^{20}$

Sol:  $q = It = n \times 2e$   
 $n = \frac{It}{2e}$   
 $= 2 \times 10^{20}$

**42.** Ans: 1, 1.2 and 1.5

Sol:  $V_A + V_B + V_C \approx 740$   
 $V_A + V_B \approx 440$   
 $V_B + V_C \approx 540$   
Solving  $V_A : V_B : V_C = 1 : 1.2 : 1.5$

**43.** Ans: The resistance of carbon decreases with the increase of temperature.

Sol: Knowledge based.

**44.** Ans:  $\pm 5\%$

Sol: Knowledge based

**45.** Ans:  $\frac{e}{2m} \lambda$

Sol:  $\frac{\mu}{L} = \frac{1}{2} \frac{e}{m}$

$$\mu = \frac{1}{2} \frac{eL}{m} = \frac{eL}{2m}$$

**46.** Ans: The radii of the wires

Sol: Knowledge based

**47.** Ans:  $\sqrt{3}$  W

Sol:  $W = mB \cos 60^\circ = mB \times \frac{1}{2}$   
 $\tau = mB \sin 60^\circ = \sqrt{3}$  W

**48.** Ans:  $\sqrt{20} \times 10^{-7}$  T

Sol:  $B = \left\{ \left[ \left( \frac{2P}{\left( \frac{d}{2} \right)^3} \right)^2 + \left( \frac{P}{\left( \frac{d}{2} \right)^3} \right)^2 \right]^{1/2} \right\} \times 10^{-7}$   
$$\sqrt{4^2 + 4} \times 10^{-7}$$
  
 $\sqrt{20} \times 10^{-7} = 2\sqrt{5} \times 10^{-7}$  T

**49.** Ans:  $1 : \sqrt{2} : 1$

Sol:  $R = \frac{mv}{qB} = \frac{\sqrt{2KE.m}}{qB}$   
 $R \propto \frac{\sqrt{m}}{q}$   
 $= 1 : \sqrt{2} : 1$

**50.** Ans:  $= 50 \mu V$

Sol:  $e = \frac{1}{2} B \lambda^2 \omega$   
 $= 50 \mu V$

**51.** Ans: 70.7 V, 70.7 mA

Sol:  $E_{rms} = \frac{E_0}{\sqrt{2}}, I_{rms} = \frac{I_0}{\sqrt{2}}$   
Each = 70.7

**52.** Ans: Eddy current

Sol: Standard result

**53.** Ans:  $\frac{1}{2} E_0 I_0 \cos \phi$

Sol: Standard result

54. Ans:  $\frac{k}{\omega}$

Sol:  $\frac{k}{\omega} = \frac{2\pi}{2\pi f} = \frac{1}{C}$

55. Ans:  $8.86 \times 10^{-12}$

Sol:  $U = \frac{1}{2} \times \frac{1}{2} \epsilon_0 E^2$   
 $= \frac{1}{4} \times 8.854 \times 10^{-12} \times (4)$   
 $= 8.86 \times 10^{-12}$

56. Ans:  $\frac{3}{4}$

Sol:  $\phi = \frac{360}{6} = 60^\circ$

$$I = I_0 \cos^2 30 = I_0 \times \frac{3}{4}$$

$$\frac{I}{I_0} = \frac{3}{4}$$

57. Ans: 40 cm

Sol:  $\frac{f_w}{f_a} = \frac{\mu - 1}{\frac{\mu_w - 1}{\mu_a}}$   
 $f_w = \frac{10 \times 0.5}{\frac{1}{8}} = 40 \text{ cm}$

58. Ans:

Sol:  $I = \frac{I_0}{2} \cdot \frac{1}{4^4} = \frac{1}{512} I_0$

59. Ans:  $\sin\theta > 8/9$

Sol:  $\mu_g = \frac{9}{8}$   
 $\sin C = \frac{1}{w \mu_g} = \frac{8}{9}$   
 $C = \sin^{-1}\left(\frac{8}{9}\right)$   
 $\theta > \sin^{-1}\frac{8}{9}$

60. Ans: 0.5 mm

Sol: Separation =  $\frac{2\lambda}{b} \times d = 0.5 \text{ mm}$

61. Ans:  $500 \text{ km s}^{-1}$

Sol:  $eV = \frac{1}{2} mv^2$   
 $v = \sqrt{\frac{e}{m} V}$   
 $= \sqrt{2 \times 1.76 \times 10^{11} \times 0.71}$   
 $= 5 \times 10^5 \text{ m s}^{-1}$   
 $= 500 \text{ km s}^{-1}$

62. Ans:  $0.4 \log_e 2$

Sol:  $R = R_0 e^{-\lambda t}$   
 $1250 = 5000 e^{-\lambda \times 5}$   
 $\lambda = 0.4 \ln 2$

63. Ans:  $\frac{r}{4}$

Sol:  $r \propto \frac{1}{p^2}$   
 $\frac{r'}{r} = \left(\frac{p}{2p}\right)^2 = \frac{1}{4}$   
 $r' = \frac{r}{4}$

64. Ans: 0.0024

Sol:  $\bar{B} = \frac{2 \times 1.115}{931}$   
 $= 0.0024 \text{ u}$

65. Ans: 0, 1, 0

Sol: x goes to zero.  
Y, Z remain unchanged.

66. Ans:  $240 \Omega$

Sol:  $P = \beta^2 \frac{R_0}{R_i}$   
 $R_i = \frac{\beta^2 R_0}{P}$   
 $= 240$

67. Ans: 3.33 mA

Sol:  $V_{R_2} = 10 \text{ V}, I_{R_2} = 6.67 \text{ mA}$   
 $I_{R_1} = \frac{5}{500} = 10 \text{ mA}$   
 $\therefore I_Z = 10 - 6.67 = 3.33 \text{ mA}$

68. Ans: Only (i) and (iii) are correct

Sol: Knowledge based.

**69.** Ans: 12.8 m

$$\text{Sol: } h = \frac{d^2}{2R} \\ = 12.8 \text{ m}$$

**70.** Ans: 50%

$$\text{Sol: } m = \frac{E_m}{E_c} = 50\%$$

**71.** Ans: Ground wave propagation is for high frequency transmission.

Sol: Knowledge based.

**72.** Ans: 8 kHz

$$\text{Sol: Band width} = 2xf_m \\ = 8 \text{ kHz}$$

**73.** Ans: II < I < III < IV

$$\text{Sol: } \begin{aligned} I &= 16 \times 1.66 \times 10^{-24} \text{ g} \\ II &= 14 \times 1.66 \times 10^{-24} \text{ g} \\ III &= 32 \times 1 \times 10^{-10} \text{ g} \\ IV &= 63 \times 1 \times 10^{-10} \text{ g} \end{aligned}$$

**74.** Ans: n = 2 to n = 1

$$\text{Sol: } \begin{aligned} \text{He}^+ (Z=2) n &= 4 \text{ to } n = 2 \\ \text{H} (Z=1) n &= \frac{4}{2} \text{ to } n = \frac{2}{2} \end{aligned}$$

**75.** Ans: O<sub>2</sub> > O<sub>2</sub><sup>-</sup>

$$\begin{array}{ll} \text{Sol: } \text{B.O} & \\ \text{C}_2 = 2 & \text{C}_2^{2-} = 3 \\ \text{B}_2^+ = 1.5 & \text{B}_2 = 2 \\ \text{Li}_2^+ = 0.5 & \text{Li}_2 = 1 \\ \text{N}_2^+ = 2.5 & \text{N}_2 = 3 \\ \text{O}_2 = 2 & \text{O}_2^- = 1.5 \end{array}$$

**76.** Ans: o-nitrophenol

Sol: Because of the proximity of the -OH and -NO<sub>2</sub> groups.

**77.** Ans: 0.67, 0.33

$$\begin{aligned} \text{Sol: } \text{C}_2\text{H}_6 + \frac{7}{2} \text{O}_2 &\rightarrow 2\text{CO}_2 + 3\text{H}_2\text{O} \\ \text{C}_2\text{H}_4 + 3\text{O}_2 &\rightarrow 2\text{CO}_2 + 2\text{H}_2\text{O} \\ \frac{7}{2}x + 3(1-x) &= \frac{10}{3} \\ x &= \frac{2}{3} \end{aligned}$$

Mole fraction of C<sub>2</sub>H<sub>6</sub> and C<sub>2</sub>H<sub>4</sub> are 0.67 and 0.33

**78.** Ans: H<sub>2</sub>O

Sol: H<sub>2</sub>O is diamagnetic

**79.** Ans: F > N > C > Be > B

Sol: F > N > C > Be > B

**80.** Ans: H<sub>2</sub>S is a reducing agent and H<sub>2</sub>O<sub>2</sub> is an oxidizing agent

Sol: H<sub>2</sub>S + H<sub>2</sub>O<sub>2</sub> → S + 2H<sub>2</sub>O  
-2 -1 0 -2  
H<sub>2</sub>S is oxidized and H<sub>2</sub>O<sub>2</sub> is reduced.

**81.** Ans: (iii) only

Sol: Carbides of Al & Be give methane with water.

**82.** Ans: NaO<sub>2</sub>

Sol: 2NaO<sub>2</sub> + 2H<sub>2</sub>O → 2NaOH + H<sub>2</sub>O<sub>2</sub> + O<sub>2</sub>

**83.** Ans: BiH<sub>3</sub>


Sol: Stability of group 15 hydride decrease down the group.

**84.** Ans: NO<sub>2</sub>

Sol: NO<sub>2</sub> dimerises on cooling to colourless N<sub>2</sub>O<sub>4</sub>

**85.** Ans: six equivalent Cr – O bonds and one Cr – O – Cr bond

Sol: Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> has the structure



There are six equivalent Cr – O bonds and one Cr – O – Cr bond

**86.** Ans: (II) and (III)

Sol: Zr<sup>+4</sup> & Hf<sup>+4</sup> is similar in size due to lanthanide contraction  
Ce<sup>+4</sup> is an oxidizing agent. La(OH)<sub>3</sub> is the most basic among lanthanide hydroxides.

**87.** Ans: 1.0 kJ

$$\begin{aligned} \text{Sol: } Q &= nC \Delta t \\ &= 2 \times 25 \times 20 \text{ J} \\ &= 1.0 \text{ kJ} \end{aligned}$$

88. Ans:  $\text{Ag}_2\text{CO}_3$  and  $\text{AgI}$

Sol:	Solubility
$\text{AgCl}$	$10^{-5}$
$\text{AgI}$	$10^{-8}$
$\text{PbCrO}_4$	$2 \times 10^{-7}$
$\text{Ag}_2\text{CO}_3$	$1.26 \times 10^{-4}$

89. Ans:  $\text{C}_6\text{H}_{12}\text{O}_6$

Sol:  $\Delta T_f = K_f \times m$   
 $0.465 = 1.86 \times \frac{1.8}{M} \times \frac{1000}{40}$   
 $M = 180$   
 $\therefore MF = \text{C}_6\text{H}_{12}\text{O}_6$

90. Ans: 16.89

Sol:  $\text{ClO}_3^- \rightarrow \text{Cl}_2$   
 $+5 \quad 0$   
Eq. mass =  $\frac{84.45}{5} = 16.89$

91. Ans:  $\frac{2}{3}$

Sol:  $9 = \left( \frac{3.24 \times 10^{-2}}{1.2 \times 10^{-3}} \right)^n$   
 $9 = (3^3)^{\frac{2}{3}}$


92. Ans: sodium stearate

Sol: Soaps and detergents are examples for associated colloids.

93. Ans: Have tetrahedral and square planar geometry respectively

Sol:  $\text{Ni}(\text{CO})_4$  is tetrahedral whereas  $[\text{Ni}(\text{CN})_4]^{2-}$  is square planar

94. Ans: 0.50



95. Ans: 72

Sol:  $\frac{5}{180} = \frac{2}{M}$   
 $M = 72$

96. Ans: 2

Sol:  $C_1 \alpha_1^2 = C_2 \alpha_2^2$

$0.1 \times (10^{-2})^2 = 0.025 \times \alpha_2^2$

$\alpha_2 = 2 \times 10^{-2}$

% = 2

97. Ans: I and IV only

Sol: For a zero order reaction rate and rate constant are independent of reactant concentration.

98. Ans:  $[\text{CoF}_6]^{3-}$

Sol:  $[\text{CoF}_6]^{3-}$  is a high spin complex containing four unpaired electrons in it.

99. Ans: -0.28 V

Sol:	$E^\circ$	$nE^\circ$
$\text{Mn}^{2+} + 2e^- \rightarrow \text{Mn}$	-1.18	-2.36 V
$\text{Mn}^{3+} + e^- \rightarrow \text{Mn}^{2+}$	1.51	1.51 V
$\text{Mn}^{3+} + 3e^- \rightarrow \text{Mn}$	-0.28	-0.85 V


100. Ans: 0.02

Sol:  $\alpha = \frac{\wedge c}{\wedge o}$   
 $= \frac{7.8}{390} = 0.02$

101. Ans:  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$

Sol: No. of moles of  $\text{AgCl}$  obtainable from 1 mole of the complex =  $\frac{430.5}{143.5} = 3$   
 $\therefore$  3 replaceable chlorines

102. Ans: 3-Methyl-2-butanone



103. Ans: Acetylene and benzene

Sol:  $\text{HC} \equiv \text{CH}$  and  $\text{C}_6\text{H}_6$  have the same empirical formula and percentage composition.

104. Ans: n-pentane > 2-methylbutane > 2, 2-dimethylpropane

Sol: As branching increases among isomeric alkanes, boiling point decreases.

105. Ans: Electrolysis

Sol: It is Kolbe's electrolytic synthesis.

**106.** Ans:  $\text{C}_6\text{H}_5 - \dot{\text{C}}\text{H} - \text{CH}_3$

Sol: It is benzylic secondary radical.

**107.** Ans: electromeric effect

Sol: Definition of electromeric effect.

**108.** Ans: Cis–2-butene and trans–2-butene

Sol: Cis–2-butene and trans–2-butene are geometrical isomers.


**109.** Ans: geometric, optical, position and functional isomerism

Sol: It can exhibit geometrical, optical, position and functional isomerism.


**110.** Ans:  $\text{CH}_3\text{Br} + \text{AgF} \rightarrow$

Sol: It is Swarts reaction


**111.** Ans: 1–bromo–3–phenylpropane



**112.** Ans: 2–methyl–2–butanol




**113.** Ans: 2–methyl–2–butanol



2-Methyl-2-butanol  
(3°alcohol)

**114.** Ans: Reimer – Tiemann reaction


Sol:



**115.** Ans: III < II < I

Sol: Order of boiling point is  $1^\circ > 2^\circ > 3^\circ$

**116.** Ans: Isopropylamine is a secondary amine



**117.** Ans: glycine and amino caproic acid

Sol: glycine and amino caproic acid are the monomers used for the preparation of Nylon–2–nylon–6.

**118.** Ans: High density polythene

Sol: HDPE is formed by the polymerization of ethane in presence of Zeigler – Natta catalyst.

**119.** Ans: cetyltrimethyl ammonium bromide

Sol: cetyltrimethyl ammonium bromide is a cationic detergent used in hair conditioners.

**120.** Ans: Food preservatives

Sol: Salts of sorbic acid and propionic acid are used as food preservatives.