

For the adiabatic approximation, the condition that $|C_m(t)| \ll 1$ implies, $|C_m(t)|^2 \rightarrow$ probability of finding the system at 'm' in a time 't' sec.

$$(a) \frac{1}{\hbar \omega_{sm}^2} \frac{\partial H'_{sm}}{\partial t} \ll 1$$

$$(b) \frac{1}{\hbar \omega_{sm}^2} \frac{\partial H'_{sm}}{\partial t} \gg 1$$

$$(c) \frac{1}{\hbar \omega_{sm}^2} \frac{\partial H'_{sm}}{\partial t} > 1$$

$$(d) \frac{1}{\hbar \omega_{sm}^2} \frac{\partial H'_{sm}}{\partial t} = 1$$

4. $J_z (J_{\pm} \psi_{jm})$ value is

$$(a) m \hbar J_{\pm} \psi_{jm}$$

$$(b) (m-1) \hbar J_{\pm} \psi_{jm}$$

$$(c) (m+1) \hbar J_{\pm} \psi_{jm-1}$$

$$(d) (m \pm 1) \hbar (J_{\pm} \psi_{jm})$$

2

4521

The matrix for Z-component of Dirac matrix α_z is

$$(a) \begin{pmatrix} 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix} \quad (b) \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$(c) \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \quad (d) \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

6. If \hat{A} be a self adjoint operator, what is the value of $\langle \phi | \hat{A} | \psi \rangle$?

7. Write down the Hamiltonian for He atom.

8. What is the percentage value of subsidiary peaks of main curve for the transition probability curves

drawn between $4 \sin^2 \left(\frac{\omega_R t}{2} \right) / \omega_R^2$ versus ω_R ?

9. Write down the matrix for L_z .

10. What is the magnetic moment for Dirac electron?

3

4521